

A note on Berwald eikonal equation

Cumali Ekici, Muradiye Çimdiker

Eskisehir Osmangazi University, Mathematics-Computer Department, Eskisehir 26480,
TURKEY

Kirklareli University, Mathematics Department, Kirklareli 39100, TURKEY

E-mail: cekici@ogu.edu.tr; muradiye.1001@hotmail.com

Abstract. In this study, firstly, we generalize Berwald map by introducing the concept of a Riemannian map. After that we find Berwald eikonal equation through using the Berwald map. The eikonal equation of geometrical optic that examining light reflects, refracts at smooth, plane interfaces is obtained for Berwald condition.

1. Introduction

Finsler manifolds can be thought as generalization of Riemannian manifolds; tangent spaces carry Minkowski norms instead of inner products and geometric objects on tangent vectors depend not only on the base but also on the fibre component. Finsler manifold have an intrinsic geometrical significance and also they have been used to model a variety of problems from dynamics, optics and relativity, cf. eg. [4] and [8].

One of the relevant figures working on that period on Finsler geometry was L. Berwald, who introduced a connection and a class of spaces sharing his name [7]. Especially, positive definite Berwald manifolds constitute the conceptually simplest and the best understood class of Finsler manifolds.

The eikonal equation is frequently solved using characteristic. This involves solving the ray equations in a combined coordinate and ray parameter phase space and then integrating the traveltime along these rays [10]. Physically rays are the trajectories along which high-frequency energy flows. A number of efficient methods for solving the ray equations have been developed [3]. Using the method of characteristics, traveltimes were computed along rays and approximate solutions of the eikonal equation could also obtained by R. L. Nowack [9].

∇f is used many areas of science such as mathematical physics and geometry. For example, the Riemannian condition $\|\nabla f\|^2 = 1$ is precisely the eikonal equation of geometrical optics. In the geometrical optical interpretation, the level sets of f are interpreted as wave fronts. The characteristics of the eikonal equation are then the solutions of the gradient flow equation for f ,

$$x' = \text{grad } f(x) \quad (1)$$

which are geodesics of M orthogonal to the level sets of f , and which are parametrized by arc length. These geodesics can be interpreted as light rays orthogonal to the wave fronts [1].

The concept of a Riemannian map was introduced by Fischer and it is shown that these maps are solutions of the eikonal equation [1]. The notion of semi-Riemannian map was stated, using the map, the solution of the eikonal equation was obtained by Garcio et al [5].



In this paper, we generalize these maps to Berwald manifolds by introducing the concept of a Riemannian map. We find Berwald eikonal equation though using the Berwald map.

2. Preliminaries

Let M be a connected, n -dimensional, C^∞ manifold and $TM = \cup_{x \in M} T_x M$ be the tangent bundle of M , where $T_x M$ is the tangent space at $x \in M$. We denote a typical point in TM by (x, y) . Set $TM_0 = TM \setminus \{0\}$, where $\{0\}$ stands for $\{(x, 0) : x \in M, 0 \in T_x M\}$. A Finsler metric on M is a function $F : TM \rightarrow [0, \infty)$ with the following properties:

i) F is C^∞ on TM_0 .

ii) At each point $x \in M$, the restriction $F_x : F|_{T_x M}$ is a Minkowski norm on $T_x M$. So the pair (M, F) is called a Finsler manifold [11].

A Finsler metric F on a manifold M is called Berwald metric if in a standard local coordinate system (x^i, y^i) in TM_0 , the Christoffel symbols $\Gamma_{jk}^i = \Gamma_{jk}^i(x)$ are functions of $x \in M$ only, in which case, $G^i = \frac{1}{2} \Gamma_{jk}^i(x) y^j y^k$ are quadratics in $y = y^i \frac{\partial}{\partial x^i} |_x$. Riemannian metrics and Minkowski metrics are trivial Berwald metrics.

The Legendre transformation $l : TM \rightarrow T^*M$ is defined as

$$l(Y) = \begin{cases} g_Y(Y, \cdot), & Y \neq 0 \\ 0, & Y = 0, \end{cases}$$

where g is the symmetric bilinear form.

Let $f : M \rightarrow \mathbb{R}$ be a smooth function on M . The gradient of f is defined as $\nabla f = l^{-1}(df)$, where $l^{-1} : T^*M \rightarrow TM$ is the inverse Legendre transformation. Thus we have

$$df(X) = g_{\nabla f}(\nabla f, X), \quad X \in TM. \quad (2)$$

[2].

Let M and M' be a Finsler manifolds with metric functions $F(x, y)$ and $F'(x', y')$. A differentiable map $f : M \rightarrow M'$ is local isometry if each of the tangent maps $f_* : T_x M \rightarrow T_{f(x)} M'$ is a vector space isomorphism which preserves the metric. A local isometry is an isometry if it is one to one and onto [6].

Let (M, F) be a Berwald space. Given any parallel vector field V along a curve σ in M . We can express the Finslerian norm of V as

$$F(V) = \sqrt{g_V(V, V)} \quad (3)$$

where $g_V = g_{ij(\sigma, V)} dx^i dx^j$ [4].

Let $f : (M, g_M) \rightarrow (N, g_N)$ be a smooth map between smooth finite dimensional Riemannian manifolds (M, g_M) and (N, g_N) . Let

$$f_* : V_x \oplus H_x \rightarrow \text{range } f_* \oplus \text{range}(f_*)^\perp$$

denote the tangent map at $x \in M$. Here $V_x = \ker f_* \subseteq T_x M$ denotes the vertical subspace of $T_x M$ and $H_x = \ker(f_*)^\perp \subseteq T_x M$ denotes the horizontal subspace of $T_x M$. If the horizontal restriction $(f_*)^h : H_x \rightarrow \text{range } f_*$ is a linear isometry between the inner product spaces $(H_x, g_M(x)|_{H_x})$ and $(\text{range } f_*, g_N(y)|_{\text{range } f_*})$, $y = f(x)$. The map is a Riemannian map if f is Riemannian at each $x \in M$ [1].

Theorem 2.1. Let $f : (M, g_M) \rightarrow (N, g_N)$ be a smooth map between smooth finite dimensional Riemannian manifolds (M, g_M) and (N, g_N) , such that M is a connected manifold.

Then $rank f$ is constant on M and the norm squared $\|f_*\|^2$ of f_* satisfies

$$\|f_*\|^2 = rank f.$$

If in the above theorem $(N, g_N) = (\mathbb{R}, 1) = \mathbb{R}$ is the real line with its Euclidean metric, and if $f : (M, g_M) \rightarrow \mathbb{R}$ is a real valued Riemannian map, then $rank f$ is constant on M equal to either zero or one. If $rank f = 0$, then f is a constant map, and if $rank f = 1$, then f satisfies the eikonal equation

$$\|f_*\|^2 = 1$$

of geometrical optics [1].

3. Berwald map and Berwald eikonal equation

Let M_1 and M_2 be differentiable, connected manifolds of dimensionals n_1 and n_2 , respectively. Let $f : (M_1, F_1) \rightarrow (M_2, F_2)$ be a map, where F_1 and F_2 are Berwald metrics. Let $f_* : T_x TM_1 \rightarrow T_y TM_2$ denotes the tangent map at $x \in M_1$, $y = f(x) \in M_2$. Then $T_x TM_1$ and $T_y TM_2$ split orthogonally as

$$T_x TM_1 = V_x TM_1 \oplus H_x TM_1$$

and

$$T_y TM_2 = range f_* \oplus range(f_*)^\perp.$$

Here

$$V_x TM_1 = \ker f_* \subseteq T_x TM_1$$

denotes the vertical subspace of $T_x TM_1$ and

$$H_x TM_1 = \ker(f_*)^\perp \subseteq T_x TM_1$$

denotes the horizontal subspace of $T_x TM_1$. Thus, we have

$$f_* : V_x TM_1 \oplus H_x TM_1 \rightarrow range f_* \oplus range(f_*)^\perp.$$

Using these expression, we can write the following definitions:

Definition 3.1. A smooth map $f : (M_1, F_1) \rightarrow (M_2, F_2)$ between smooth finite dimensional (M_1, F_1) and (M_2, F_2) is Berwald at $x \in M_1$ if the horizontal restriction

$$(f_*)^h : H_x TM_1 \rightarrow range f_*$$

is a linear isometry between $(H_x TM_1, F_1(x)|_{H_x})$ and $(range f_*, F_2(y)|_{range f_*})$, $y = f(x)$. The map is a Berwald map if f is Berwald at each $x \in M_1$.

Definition 3.2. Let $f : (M_1, F_1) \rightarrow (M_2, F_2)$ be a map. Then the square norm of f_* is the map $\|f_*\|^2 : M_1 \rightarrow \mathbb{R}^+$ defined by $\|f_*\|^2(p_1) = \|f_{*p_1}\|^2$.

Note that, since $\|f_*\|^2 = \sum_{i=1}^{n_1} F_1^2(X_i)F_2^2(f_*(X_i))$, where $\{X_1, \dots, X_{n_1}\}$ is an orthonormal local frame of TTM_1 .

Let us compute the square norm of a map $f : (M, F) \rightarrow (\mathbb{R}^+, dt \otimes dt)$. For this, let $\{X_1, \dots, X_n\}$ be an orthonormal local frame. We have

$$\|f_*\|^2 = \sum_{i=1}^n F^2(X_i)(dt \otimes dt)(f_*X_i, f_*X_i)$$

From here, using the expression (2), then we find

$$\begin{aligned} \|f_*\|^2 &= \sum_{i=1}^n F^2(X_i)(g_{\nabla f}(\nabla f, X_i))^2(dt \otimes dt) \left(\frac{d}{dt} \circ f, \frac{d}{dt} \circ f \right) \\ &= g_{\nabla f}(\nabla f, \nabla f) \end{aligned}$$

From the equation (3), we have $\|f_*\|^2 = F^2(\nabla f)$. The generalized eikonal equations for $f : (M, F) \rightarrow (\mathbb{R}^+, dt \otimes dt)$ become $\|f_*\| = F(\nabla f) = 1$. This extension enables a fast calculation of geodesic paths.

Remark 3.1. According to the Theorem 2.1, for Berwald, this case is a geometrical optical condition which it describes light propagation in terms of rays.

Lemma 3.1. Let $f : (M, F) \rightarrow (\mathbb{R}^+, dt \otimes dt)$ be a map. Then

$$\nabla(F^2(\nabla f)) = 2\nabla_{\nabla f}\nabla f,$$

where ∇f and $\nabla(F^2(\nabla f))$ denote the gradients of f and $F^2(\nabla f)$ on Berwald manifolds.

Proof. If $X \in \Gamma TM$, then we have

$$g(\nabla(F^2(\nabla f)), X) = 2g_{\nabla f}(\nabla_X \nabla f, \nabla f)$$

Since $h_f(X) = \nabla_X \nabla f$, we find

$$g(\nabla(F^2(\nabla f)), X) = 2g_{\nabla f}(\nabla_{\nabla f} \nabla f, X).$$

Hence, this completes the proof.

Proposition 3.1. Let $f : (M, F) \rightarrow (\mathbb{R}^+, dt \otimes dt)$ be a map. If f satisfies a Berwald eikonal equation $F^2(\nabla f) = 1$, then ∇f is a geodesic vector field on (M, F) .

Proof. For Berwald eikonal equation, we have $F^2(\nabla f) = 1$. From Lemma 3.1., we find

$$\nabla(F^2(\nabla f)) = 2\nabla_{\nabla f}\nabla f = 0. \tag{4}$$

Therefore, ∇f is a geodesic vector field on (M, F) . Namely, integral curves of the gradient flow of f are geodesics of the Berwald manifolds.

Proposition 3.2. Let $f : (M, F) \rightarrow (\mathbb{R}^+, dt \otimes dt)$ be a map. Then

i) The rank of f is constant on M .

ii) $F^2(\nabla f)$ is constant on M .

Proof. The rank of f is constant on M if and only if for every $p \in M$ either $\nabla f(p) = 0$ or $\nabla f(p) \neq 0$. Here, we will only notice that $\nabla f(p) \neq 0$ for every $p \in M$.

i) Let $p, q \in M$ and $\gamma : [a, b] \rightarrow M$ be a curve with $\gamma(a) = p, \gamma(b) = q$. Since ∇f is a parallel vector field on (M, F) , $(\nabla f) \circ \gamma$ is parallel vector field along γ . By the uniqueness of parallel vector fields along a curve with respect to initial condition, $\nabla f(p) \neq 0 \Leftrightarrow \nabla f(q) \neq 0$. Thus, we have $\nabla f(p) \neq 0$ for every $p \in M$.

ii) According to the expression (4), $F^2(\nabla f)$ is constant on M .

4. Conclusion

In this work, we introduce Berwald map and study to explain the eikonal equation of geometrical optic for Berwald condition. In future work, we will consider affine solution of Berwald eikonal equation. First of all, the physical implications of affine solution of Berwald eikonal equation should be investigated. Last, an obvious next step would be to show how to use this theorem where generalizations of Berwald metric are employed in order to construct the eikonal equation in pseudo-Finsler space.

5. References

- [1] A. E. Fischer, 1992, Riemannian Map Between Riemannian Manifolds, *Contemporary Math*, **182**, 342.
- [2] B. Y. Wu, 2013, Comparison Theorems in Finsler geometry with Weighted Curvature Bounds and Related Results, *J. Korean Math. Soc.*, **52**(3), 603-624.
- [3] B. Julian and D. Gubbins, 1977, Three Dimensional Seismic Ray Tracing, *Geophys. J. Int.*, **43**, 95-113.
- [4] D. Bao, S. S. Chern and Z. Shen, 2000, An Introduction to Riemann- Finsler Geometry, Springer-Verlag, Newyork.
- [5] E. Garcio and D. N. Küpeli, 1999, Semi- Riemannian Maps and Their Applications, Kluwer Academic Publishers, London.
- [6] J. K. Beem, 1971, Motions in Two Dimensional Indefinite Finsler Spaces, *Indiana University Mathematics Journal*, **21**(6).
- [7] L. Berwald, 1925, Untersuchung der Krümmung Allgemeiner Metrischen Raume auf Grund des in Ihnen Herrschenden Parallelismus, *Math. Zeit*, **25**, 40-73.
- [8] P. L. Antonelli, R. S. Ingarden and M. Matsumoto, 1993, The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology, Kluwer Academic Publishers.
- [9] R. L. Nowack, 1992, Wavefronts and Solutions of the Eikonal Equation, *Geophys. J. Int.*, **110**, 55-62.
- [10] V. Cerveny, I. A. Molotkov and I. Psencik, 1977, Ray Method in Seismology, Praha, Univ. Karlova.
- [11] Z. Shen, 2004, Landsberg Curvature, S -Curvature and Riemann Curvature, In: A Sampler of Finsler Geometry, MSRI Publications, **50**, Cambridge Univ. Press.